

Documents

Shibl, M.F.^a, Elroby, S.A.K.^b, Hilal, R.H.^a

Solvent and substituent effects on the electronic structures of triazoles: Computational study
(2011) *Molecular Simulation*, 37 (1), pp. 11-17.

^a Chemistry Department, Faculty of Science, Cairo University, Cairo, Egypt

^b Chemistry Department, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia

Abstract

The tautomerisation reaction of 1,2,3-triazole and some of its derivatives has been studied and the transition states were located. The analysis of results indicates that 2H-triazole is more stable than the 1H-tautomer in all studied compounds in gas phase and in solution. The proton detachment energies and the geometries of the compounds studied were analysed using density functional theory (DFT) and Hartree-Fock (HF) methods in gas phase and solution. Deprotonation takes place in all cases studied by the detachment of N-H proton. The solvent effect on the stability the tautomeric and anion forms of studied compounds has been examined using B3LYP/6-31G* level of theory by applying the polarisable continuum model. © 2011 Taylor & Francis.

Author Keywords

activation energy; detachment energies (PDE); PCM; tautomerism; triazole

Document Type: Article

Source: Scopus

About Scopus

[What is Scopus](#)
[Content coverage](#)
[What do users think](#)
[Latest](#)
[Tutorials](#)

Contact and Support

[Contact and support](#)
[Live Chat](#)

About Elsevier

[About Elsevier](#)
[About SciVerse](#)
[About SciVal](#)
[Terms and Conditions](#)
[Privacy Policy](#)

